Search

**Ophthalmology and Visual Sciences** 




# Central posterior synechiae (CPS) in the setting of tubulointerstitial nephritis and uveitis syndrome (TINU)


Category(ies): Uveitis

Contributor: <u>Jesse Vislisel, MD</u> Photographer: Toni Venckus, CRA



Central posterior synechiae (CPS) are adhesions between the iris at the pupillary margin and anterior lens capsule that can occur in settings of intraocular inflammation. This patient has panuveitis related to tubulointerstitial nephritis and uveitis syndrome (TINU). As seen here, the CPS are very apparent after pharmacologic dilation as they mechanically restrict the pupil from expanding.





# **Image Permissions:**



Ophthalmic Atlas Images by <u>EyeRounds.org</u>, <u>The University of Iowa</u> are licensed under a <u>Creative Commons Attribution-NonCommercial-</u> NoDerivs 3.0 Unported License.

#### Address

University of Iowa Roy J. and Lucille A. Carver College of Medicine Department of Ophthalmology and Visual Sciences

### Legal

Copyright © 2019 The University of Iowa. All Rights Reserved Report an issue with this page Web Privacy Policy | Nondiscrimination Statement

# **Related Links**

Cataract Surgery for Greenhorns EyeTransillumination Gonioscopy.org Iowa Glaucoma Curriculum Iowa Wet Lab Patient Information

# **EyeRounds Social** Media

Follow







200 Hawkins Drive Iowa City, IA 52242 Stone Rounds The Best Hits Bookshelf Contact Us Submit a Suggestion

Support Us