

Traction retinal detachment secondary to proliferative diabetic retinopathy

Category(ies): Retina, Vitreous

Contributor: [Elizabeth Gauger, MD](#)

Photographer: Stefani Karakas, CRA



44-year-old female with a 25 year history of poorly-controlled insulin dependent diabetes mellitus. She noticed a decrease in vision in the left eye over several months.



This photo shows a tractional retinal detachment secondary to proliferative diabetic retinopathy. There are fibrovascular bands encircling the arcades that are fovea (macula) threatening. Please note the neovascularization of the disc (NVD) originating from the inferonasal edge of the optic disc.

[Enlarge](#) [Download](#)

Image Permissions:



Ophthalmic Atlas Images by [EyeRounds.org](#), [The University of Iowa](#) are licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike](#)

Address

University of Iowa
Roy J. and Lucille A. Carver College
of Medicine
Department of Ophthalmology and
Visual Sciences
200 Hawkins Drive
Iowa City, IA 52242

[Support Us](#)

Legal

Copyright © 2019 The University of
Iowa. All Rights Reserved
[Report an issue with this page](#)
[Web Privacy Policy](#) |
[Nondiscrimination Statement](#)

Related Links

[Cataract Surgery for Greenhorns](#)
[EyeTransillumination](#)
[Gonioscopy.org](#)
[Iowa Glaucoma Curriculum](#)
[Iowa Wet Lab](#)
[Patient Information](#)
[Stone Rounds](#)
[The Best Hits Bookshelf](#)

EyeRounds Social Media

Follow



[Receive notification of new cases,
sign up here](#)
[Contact Us](#)
[Submit a Suggestion](#)